

16 *Years*

Previous Years Solved Papers

Civil Services Main Examination

(2009-2024)

Mathematics Paper-II

Topicwise Presentation





MADE EASY Publications Pvt. Ltd.

Corporate Office: 44-A/4, Kalu Sarai (Near Hauz Khas Metro Station), New Delhi-110016

E-mail: infomep@madeeasy.in

Contact: 9021300500

Visit us at: www.madeeasypublications.org

Civil Services Main Examination Previous Solved Papers : Mathematics Paper-II

© Copyright, by MADE EASY Publications Pvt. Ltd.

All rights are reserved. No part of this publication may be reproduced, stored in or introduced into a retrieval system, or transmitted in any form or by any means (electronic, mechanical, photo-copying, recording or otherwise), without the prior written permission of the above mentioned publisher of this book.

First Edition: 2020

Second Edition: 2021

Third Edition: 2022

Fourth Edition: 2023

Fifth Edition: 2024

Sixth Edition: 2025

Preface

Civil Service is considered as the most prestigious job in India and it has become a preferred destination by all engineers. In order to reach this estimable position every aspirant has to take arduous journey of Civil Services Examination (CSE). Focused approach and strong determination are the pre-requisites for this journey. Besides this, a good book also comes in the list of essential commodity of this odyssey.



I feel extremely glad to launch the fourth edition of such a book which will not only make CSE plain sailing, but also with 100% clarity in concepts.

MADE EASY team has prepared this book with utmost care and thorough study of all previous years papers of CSE. The book aims to provide complete solution to all previous years questions with accuracy.

I would like to acknowledge efforts of entire MADE EASY team who worked day and night to solve previous years papers in a limited time frame and I hope this book will prove to be an essential tool to succeed in competitive exams and my desire to serve student fraternity by providing best study material and quality guidance will get accomplished.

With Best Wishes

B. Singh (Ex. IES)

CMD, MADE EASY Group

Previous Years Solved Papers of
Civil Services Main Examination

Mathematics : Paper-II

C O N T E N T S

Sl.	TOPIC	PAGE No.
Unit-1	Modern Algebra	1 - 45
	1. Groups.....	1
	2. Rings	23
	3. Fields.....	38
Unit-2	Real Analysis	46 - 101
	1. Real Number System	46
	2. Sequences	46
	3. Series	54
	4. Functions	66
	5. Riemann Integral.....	82
	6. Improper Integral.....	89
	7. Functions of Several (Two or Three) Variables	92
Unit-3	Complex Analysis.....	102 - 146
	1. Analytic Functions	102
	2. Taylor's Series	117
	3. Singularity	122
	4. Laurant's Series	124
	5. Cauchy's Residue Theorem.....	132
	6. Contour Integration	137

Unit-4	Linear Programming Problems.....	147 - 199
	1. Basic Feasible Solutions.....	147
	2. Optimal Solutions.....	150
	3. Graphical Method of Solution.....	153
	4. Simplex Method of Solution.....	159
	5. Duality.....	178
	6. Transportation Problems.....	182
	7. Assignment Problems.....	191
Unit-5	Partial Differential Equations	200 - 289
	1. Formulation of P.D.E.....	200
	2. Solution of Quasilinear Partial Differential Equations of 1st Order, Lagrange's Auxiliary Equations and Charpit's Auxiliary Equations	208
	3. Cauchy's Method of Characteristics	220
	4. Linear P.D.E. of Second Order with Constant Coefficient.....	225
	5. Canonical Form.....	237
	6. Application of PDE.....	255
Unit-6	Numerical Analysis and Computer Programming	290 - 353
	1. Solution of Algebraic and Transcendental Equation of One Variable.....	290
	2. Solutions of System of Linear Equations.....	296
	3. Interpolation.....	310
	4. Numerical Integration.....	321
	5. Ordinary Differential Equation.....	330
	6. Computer Programming.....	334
Unit-7	Mechanics and Fluid Dynamics	354 - 430
	1. Generalised Coordinate.....	354
	2. Moment of Inertia.....	375
	3. Motion of Rigid Bodies in Two Dimensions.....	386
	4. Equation of Continuity.....	389
	5. Euler's Equation of Motion for Inviscid Flow	402
	6. Source and Sink.....	410
	7. Vortex Motion	416
	8. Navier-Stoke's Equation for Viscous Flow.....	427



1. Groups

1.1 If \mathbb{R} is the set of real number and \mathbb{R}_+ is the set of positive real numbers, show that \mathbb{R} under addition $(\mathbb{R}, +)$ and \mathbb{R}_+ under multiplication (\mathbb{R}_+, \cdot) are isomorphic. Similarly, if \mathbb{Q} is the set of rational numbers and \mathbb{Q}_+ the set of positive rational numbers, are $(\mathbb{Q}, +)$ and (\mathbb{Q}_+, \cdot) isomorphic? Justify your answer.

(2009 : 4+8 =12 Marks)

Solution:

Let \mathbb{R} be the set of real number and \mathbb{R}_+ be the set of positive real number.

We have to show

$$(\mathbb{R}, +) \cong (\mathbb{R}_+, \cdot)$$

Define $f: \mathbb{R} \rightarrow \mathbb{R}_+$ as

$$f(x) = a^x; \text{ where } a > 0.$$

We will show f is one-one.

Consider,

$$\begin{aligned} \ker f &= \{x \in \mathbb{R} \mid f(x) = 1\} \\ &= \{x \in \mathbb{R} \mid a^x = 1\} \\ &= \{x \in \mathbb{R} \mid x = \log_a 1\} \\ &= \{x \in \mathbb{R} \mid x = 0\} \\ &= \{0\} \end{aligned}$$

$\therefore f$ is 1-1.

We will show f is homomorphism.

Let $x, y \in \mathbb{R}$

Consider

$$\begin{aligned} f(x+y) &= a^{x+y} \\ &= a^x \cdot a^y = f(x) \cdot f(y) \end{aligned}$$

$\therefore f$ is homomorphism. We will show f is onto, i.e., we have to find for any positive real number 'y' some real number x such that

$$f(x) = y$$

i.e.,

$$a^x = y$$

As

$$a^x = y$$

On taking log both sides

\Rightarrow

$$x = \log_a y$$

\therefore

$$f(x) = y$$

Hence, f is onto.

\therefore

$$(\mathbb{R}, +) \cong (\mathbb{R}_+, \cdot)$$

Let \mathbb{Q} be the set of rational numbers and \mathbb{Q}_+ be the set of positive rational number.

If f is homomorphism from \mathbb{Q} to \mathbb{Q}_+ , then

$$f(x, y) = f(x)f(y) \quad \forall x, y \in Q$$

And if image of 1 is known then the image of every element will be known.

$$\therefore f(x) = a^x \text{ where } a = f(1)$$

$$\text{If } a = 1, \quad f(x) = 1$$

$\therefore f$ is trivial homomorphism.

$$\text{If } a \neq 1, \quad f(x) = 1$$

$$\text{then } f(x) = a^x \in Q_+ \quad \forall x \in Q$$

which is a contradiction.

Hence, only trivial homomorphism is possible.

$$\therefore (Q, +) \not\cong (Q_+, \cdot)$$

1.2 Determine the number of homomorphisms from the additive group Z_{15} to the additive group Z_{10} . (Z_n is the cyclic group of order n).

(2009 : 12 Marks)

Solution:

Let $\phi : Z_{15} \rightarrow Z_{10}$ be a homomorphism.

As Z_{15} is a cyclic group of order 15.

$$Z_{15} = \langle 1 \rangle$$

Under homomorphism, if element 1 will be mapped then remaining elements will get mapped themselves ($\because G$ is cyclic)

$$\text{Suppose, } \phi(1) = x$$

As we know, if f is homomorphism from G to G' then $O(f(a)) \mid O(a)$ where $a \in G$.

As $\phi(1) = x$.

$$\therefore O(x) \mid O(1) = 15$$

And order of element divides order of group

$$\therefore O(x) \mid 10 \text{ As } O(x) \mid 15$$

$$O(x) \mid 15 \Rightarrow O(x) \mid \text{g.c.d.}(15, 10) = 5$$

$$\therefore O(x) = 1 \text{ or } O(x) = 5$$

If $O(x) = 1$. Then it is trivial homomorphism. And if $O(x) = 5$.

Note : In Z_n , number of elements of order $k = \phi(k)$; provided $k \mid n$.

$$\therefore \text{In } Z_{10}, \text{ number of elements of order } 5 = \phi(5) = 4.$$

\therefore We have 4 possibilities for x .

$$\text{Total number of homomorphism} = 4 + 1 = 5.$$

1.3 Show that the alternating group on four letters A_4 has no subgroup of order 6.

(2009 : 15 Marks)

Solution:

Consider the alternating group A_4 .

$$\sigma(A_4) = \frac{\sigma(S_4)}{2} = \frac{12}{2} = 6$$

We show although $6 \mid 12$, A_4 has no subgroup of order 6. Suppose H is a subgroup of A_4 and $\sigma(H) = 6$.

By previous problem the number of distinct 3-cycles in S_4 is

$$\frac{1}{3} \cdot \frac{4!}{(4-3)!} = \frac{4 \cdot 3 \cdot 2 \cdot 1}{3 \cdot 1} = 8$$

Again, as each 3-cycle will be even permutation all these 3-cycles are in A_4 . Obviously then, at least one 3-cycle, say s does not belong to H ($\sigma(H) = 6$).

Now, $\sigma \in H \Rightarrow \sigma^2 \in H$, because if $\sigma^2 \in H$.

Then, $\sigma^4 \in H$

$\Rightarrow \sigma \in H$

As $\sigma^3 = I$ as $\sigma(\sigma) = 3$

Let $K = \langle \sigma \rangle = \{I, \sigma, \sigma^2\}$ then, $\sigma K = 3 (= \sigma(\sigma))$

and $H \cap K = \{1\}$ ($\sigma, \sigma^2 \notin H$)

$$\Rightarrow \sigma(HK) = \frac{\sigma(H) \cdot \sigma(K)}{\sigma(H \cap K)} = \frac{6 \cdot 3}{1} = 18,$$

not possible as $HK \subseteq A_4$ and $\sigma(A_4) = 12$.

1.4 Let $G = \mathbb{R} - \{-1\}$ be the set of all real numbers omitting -1 . Define the binary relation $*$ on G by $a*b = a + b + ab$. Show $(G, *)$ is a group and it is abelian.

(2010 : 12 Marks)

Solution:

Given : Binary relation $*$ as

$$a*b = a + b + ab, \text{ where } a, b \in G.$$

Closure : Let $a, b \in G$

$$\therefore a*b = a + b + ab$$

if $a + b + ab = -1$, then

$$a + b + ab + 1 = 0$$

$$\Rightarrow (1 + a) + b(1 + a) = 0$$

$$\Rightarrow (1 + a) + (1 + b) = 0$$

$$\Rightarrow \text{either } 1 + a = 0 \Rightarrow a = -1$$

$$\text{or } 1 + b = 0 \Rightarrow b = -1$$

\therefore both $a, b \in G$

$$\therefore a \neq -1 \text{ and } b \neq -1$$

$$\therefore a + b + ab \neq -1 \text{ for any } a, b \in G$$

$$\therefore a + b + ab \neq -1 \text{ for any } a, b \in G$$

$$\therefore a*b \in G$$

So, closure is satisfied.

Associative : Let $a, b, c \in G$

$$\begin{aligned} \therefore (a*b)*c &= (a + b + ab)*c \\ &= a + b + ab + c + (a + b + ab)c \\ &= a + b + ab + c + ac + bc + abc \\ &= a + b + c + ab + ac + bc + abc \end{aligned}$$

$$\begin{aligned} \text{Also, } a*(b*c) &= a*(b + c + bc) \\ &= a + b + c + bc + a(b + c + bc) \\ &= a + b + c + ab + bc + ac + abc \end{aligned}$$

$$\text{as } (a*b)*c = a*(b*c)$$

\therefore Associative property is satisfied.

Identity :

$$\text{Let } a*b = a = a + b + ab$$

$$\begin{aligned} \Rightarrow a + b + ab &= a \\ \Rightarrow b(1 + a) &= 0 \\ \text{as } a &\neq -1 \Rightarrow b = 0 \end{aligned}$$

$\therefore b = 0$ is an identity and as $O \in G$.

\therefore identity exists.

Inverse :

Let $a * b = 0 = a + b + ab$

$$\Rightarrow a(1 + b) = -b \Rightarrow a = \frac{-b}{1+b} \quad (b \neq -1)$$

Also, $\frac{-b}{1+b} \in G$

\therefore Inverse exists.

As closure, associative property, identity, inverse conditions are satisfied. $\therefore (G, *)$ is a group.

Now, $a * b = a + b + ab$
 $b * a = b + a + ba = a + b + ab$

as $a * b = b * a$

$\therefore (G, *)$ is abelian.

- 1.5 Show that a cyclic group of order 6 is isomorphic to the product of a cyclic group of order 2 and a cyclic group of order 3. Can you generalize this? Justify.

(2010 : 12 Marks)

Solution:

Let Z_6 is cyclic group of order 6
 Z_2 is cyclic group of order 2
 Z_3 is cyclic group of order 3

Now, we know that

$$Z_m \times Z_n \cong Z_{mn}$$

when m and n are co-prime.

Here, $m = 2, n = 3$

$\therefore Z_2 \times Z_3 \cong Z_6$

- 1.6 Let (\mathbb{R}^*, \cdot) be the multiplicative group of non-zero real and $(GL(n, \mathbb{R}), X)$ be the multiplicative group of $n \times n$ non-singular real matrices. Show that the quotient group $GL(n, \mathbb{R}) / SL(n, \mathbb{R})$ and (\mathbb{R}^*, \cdot) are isomorphic where

$$SL(n, \mathbb{R}) = \{A \in GL(n, \mathbb{R}) / \det A = 1\}$$

What is the center of $GL(n, \mathbb{R})$?

(2010 : 15 Marks)

Solution:

Given, $(GL(n, \mathbb{R}), X)$ is multiplicative group of real matrices.

Let $f: GL(n, \mathbb{R}) \rightarrow (\mathbb{R}^*, \cdot)$ be a no morphism where for any $A \in G$.

$$\begin{aligned} \phi(A) &= |A| \\ \therefore \phi(A \times B) &= |AB| = |A| |B| \end{aligned}$$

Now, let $\text{Ker}(\phi)$ contains matrices such it $B \in \text{Ker}(\phi)$

then $\phi(AB) = A$

$\therefore |B| = 1$

$\therefore \text{Ker}(\phi)$ contains matrices such that $|B| = 1$ if $B \in \text{Ker}(\phi)$.

\therefore By first fundamental theorem of homomorphism,

$$\frac{GL}{\text{Ker}(\phi)} \cong (\mathbb{R}^*, \cdot)$$

Given, $SL(n, \mathbb{R}) = \{A \in GL(n, \mathbb{R}) / |A| = 1\}$

i.e., $SL(n, \mathbb{R})$ is $\text{Ker}(\phi)$.

$$\therefore \frac{GL(n, \mathbb{R})}{SL(n, \mathbb{R})} \cong (\mathbb{R}^*, \cdot)$$

Now, let z be the center of $GL(n, \mathbb{R})$.

for $x \in z$ where $X = \begin{bmatrix} p & q \\ r & s \end{bmatrix}$

$\forall A \in GL(n, \mathbb{R}), AX = XA$

Let $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ where $A \in GL(n, \mathbb{R})$

Now, $AX = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} p & q \\ r & s \end{bmatrix} = \begin{bmatrix} ap+br & aq+bs \\ cp+dr & cq+ds \end{bmatrix}$

$XA = \begin{bmatrix} p & q \\ r & s \end{bmatrix} \begin{bmatrix} a & b \\ c & d \end{bmatrix} = \begin{bmatrix} ap+qc & pb+qd \\ ar+cs & br+ds \end{bmatrix}$

\therefore for $XA = AX$

$$\begin{bmatrix} ap+br & aq+bs \\ cp+dr & cq+ds \end{bmatrix} = \begin{bmatrix} ap+qc & pb+qd \\ ar+cs & br+ds \end{bmatrix}$$

Holds true when $p = q = r = s$

or $p = s, q = r = 0$

$$\therefore z = \left\{ \begin{bmatrix} p & p \\ p & p \end{bmatrix}, \begin{bmatrix} p & 0 \\ 0 & p \end{bmatrix} \mid p \in R \right\}$$

1.7 Show that the set

$$G = \{f_1, f_2, f_3, f_4, f_5, f_6\}$$

of six transformations on the set of Complex numbers defined by

$$f_1(z) = z, f_2(z) = 1 - z, f_3(z) = \frac{z}{(z-1)},$$

$$f_4(z) = \frac{1}{z}, f_5(z) = \frac{1}{(1-z)} \text{ and } f_6(z) = \frac{(z-1)}{z}$$

is a non-abelian group of order 6 w.r.t. composition of mappings.

(2011 : 12 Marks)

Solution:

Let us construct the following table of the elements of G w.r.t. composition of mappings :

	f_1	f_2	f_3	f_4	f_5	f_6
f_1	z	$1-z$	$\frac{z}{z-1}$	$\frac{1}{z}$	$\frac{1}{1-z}$	$\frac{z-1}{z}$
f_2	$1-z$	z	$\frac{1}{1-z}$	$\frac{z-1}{z}$	$\frac{z}{z-1}$	$\frac{1}{z}$
f_3	$\frac{z}{z-1}$	$\frac{z-1}{z}$	z	$\frac{1}{1-z}$	$\frac{1}{z}$	$1-z$
f_4	$\frac{1}{z}$	$\frac{1}{1-z}$	$\frac{z-1}{z}$	z	$1-z$	$\frac{z}{z-1}$
f_5	$\frac{1}{1-z}$	$\frac{1}{z}$	$1-z$	$\frac{z}{z-1}$	$\frac{z-1}{z}$	z
f_6	$\frac{z-1}{z}$	$\frac{z}{z-1}$	$\frac{1}{z}$	$1-z$	z	$\frac{1}{1-z}$

From the table it is clear that

$$(f_i \circ f_i)(z) = (f_i \circ f_i)(z) \quad \forall z \in C, 1 \leq i \leq 6$$

$\therefore f_i$ is the identity of G .

Also,

$$f_1^{-1} = f_1; f_2^{-1} = f_2; f_3^{-1} = f_3; f_4^{-1} = f_4; f_5^{-1} = f_5; f_6^{-1} = f_6$$

$\Rightarrow G$ is a group.

But

$$(f_2 \circ f_3)(z) = \frac{1}{1-z}$$

and

$$(f_3 \circ f_2)(z) = \frac{z-1}{z}$$

$\therefore G$ is a non-abelian group.

1.8 (i) Prove that a group of Prime order is abelian.

(2011 : 6 Marks)

(ii) How many generators are there of the cyclic group (G, \cdot) of order 8?

(2011 : 6 Marks)

Solution:

(i) Let $O(G) = P$, where P is a prime number.

Let $a \in h$ be any element.

Since order of an element divides the order of the group.

$$\therefore O(a) \mid O(G) = P$$

$$\Rightarrow O(a) = 1 \text{ or } O(a) = P$$

If $a \neq e$, then

$$O(a) = P$$

Let

$$H = \langle a \rangle$$

\Rightarrow

$$O(H) = O(a) = P \text{ and } H \subseteq G$$

\therefore

$$H = G = \langle a \rangle$$

$\Rightarrow G$ is cyclic.

$\Rightarrow G$ is abelian (as every cyclic group is abelian).

(ii) Let G be a cyclic group of order 8.

$\therefore \exists$ an element $a \in G$ such that $O(a) = 8$

We know that,

$$O(a^n) = \frac{O(a)}{(O(a), n)}$$

\therefore

$$O(a^n) = O(a) \Rightarrow (O(a), n) = 1$$

Number of elements of G whose order is co-prime to

$$\begin{aligned} O(a) &= \phi(O(a)) \\ &= \phi(8) = \phi(2^3) \\ &= 4 \end{aligned} \quad (\because \phi(P^n) = P^n - P^{n-1})$$

\therefore Number of generators of cyclic group of order 8 = 4.

1.9 Give an example of a group G in which every proper subgroup is cyclic but the group itself is not cyclic.

(2011 : 15 Marks)

Solution:

Let $G = Q_8 = \{\pm 1, \pm i, \pm j, \pm k\}$

Define product on G by usual multiplication together with

$$\begin{aligned} i^2 &= j^2 = k^2 = -1 \\ ij &= -ji = k \\ jk &= -kj = i \\ ki &= -ik = j \end{aligned}$$

Then G forms a group.

But G is not abelian as $ij \neq ji$.

$\Rightarrow G$ can not be cyclic as every cyclic group is abelian.

Let

$$\begin{aligned} H_1 &= [1] = \langle 1 \rangle \\ H_2 &= [\pm 1] = \langle -1 \rangle \\ H_3 &= [\pm 1, \pm i] = \langle i \rangle = \langle -i \rangle \\ H_4 &= [\pm 1, \pm j] = \langle j \rangle = \langle -j \rangle \\ H_5 &= [\pm 1, \pm k] = \langle k \rangle = \langle -k \rangle \end{aligned}$$

Thus, H_1 to H_5 are proper subgroups of G and all there are cyclic. Hence the result.

1.10 Let a and b be elements of a group, with $a^2 = e$, $b^6 = e$ and $ab = b^4a$. Find the order of ab , and express its inverse in each of the forms $a^m b^n$ and $b^m a^n$.

(2011 : 20 Marks)

Solution:

Given : $a^2 = e, b^6 = e$ and $ab = b^4a$

$\therefore ab = b^4a$

$\Rightarrow aba^{-1} = b^4aa^{-1} = b^4e = b^4$

Again, $b^8 = b^4 \cdot b^4$

$$= (aba^{-1})(aba^{-1}) = ab^2a^{-1}$$

$\Rightarrow b^{16} = b^8 \cdot b^8 = (ab^2a^{-1})(ab^2a^{-1})$

$$= ab^4a^{-1}$$

$$= a(aba^{-1})a^{-1} \quad (\because b^4 = aba^{-1})$$

$$= a^2ba^{-2}$$

$$= a^2b(a^2)^{-1}$$

$$= ebe = b$$

$\Rightarrow b^{16} = b \Rightarrow b^{15} = e$

$\Rightarrow O(b) = 1, 3, 5$ or 15 ... (i)

Also, given $b^6 = e$

$\Rightarrow O(b) = 1, 2, 3$ or 6 ... (ii)

\therefore From (i) and (ii)

$$O(b) = 3$$

And $ab = b^4a = b^3ba = eba = ba$

Also, we know that, for $a, b \in G$ such that $ab = ba$ and $(O(a), O(b)) = 1$

$$\text{then } O(ab) = O(a) \cdot O(b)$$

$$\therefore O(ab) = 2 \times 3 = 6$$

$$\Rightarrow (ab)^6 = e$$

$$\Rightarrow (ab)^5 = (ab)^{-1}$$

$$\Rightarrow a^5 b^5 = (ab)^{-1}$$

$$\text{Similarly, as } ab = ba, (ab)^{-1} = b^5 a^5$$

1.11 How many elements of order 2 are there in the group of order 16 generated by a and b such that the order of a is 8, the order of b is 2 and $bab^{-1} = a^{-1}$.

(2012 : 12 Marks)

Solution:

Let G be the given group of order 16.

$$\text{Then, } O(G) = 16$$

$$O(a) = 8$$

$$O(b) = 2$$

$$\text{and } bab^{-1} = a^{-1} \quad \dots(i)$$

From (i), we have

$$b(bab^{-1}) = b(a^{-1})$$

$$\Rightarrow b^2 ab^{-1} = ba^{-1}$$

$$\Rightarrow ab^{-1} = ba^{-1} \quad (\because O(b) = 2 \Rightarrow b^2 = e)$$

$$\Rightarrow ab = ba^{-1} \quad (\because b^2 = e \Rightarrow b = b^{-1})$$

and h is generated by a and b , where $O(a) = 8$, $O(b) = 2$, $bab^{-1} = a^{-1}$.

$\therefore G = D_8$, dihedral group of order 8.

$$\text{Again, let } H = \langle a \rangle$$

$$\text{Since, } O(a) = 8$$

$\therefore H$ is a cyclic group of order 8. Also, in a finite cyclic group, for each divisor $\frac{k}{n} = O(G)$, there exist exactly

$\phi(k)$ elements of order k .

$[\phi(n) = \text{positive integers upto a given integer } n \text{ that are relatively prime to } n].$

As $2 \mid 8$, $\therefore \phi(2)$ elements of order 2, i.e., 1 element of order 2 ($\because \phi(2) = 1$)

Also, D_8 consists of 8 rotations and 8 reflections.

Since, order of each reflection is 2.

\therefore Number of elements of order 2 = 8 + 1 = 9.

1.12 How many conjugacy classes does the permutation group S_5 of permutations 5 numbers have? Write down one element in each class (preferably in terms of cycles).

(2012 : 15 Marks)

Solution:

We know that two permutations are said to be conjugate if they have same cyclic decomposition.

There are 7 cyclic decompositions which are :

- | | | |
|----|---------------------|---------------------------|
| 1. | $1 + 1 + 1 + 1 + 1$ | [1 (5 times)] |
| 2. | $1 + 1 + 1 + 2$ | [2 (1 time), 1 (3 times)] |
| 3. | $1 + 1 + 3$ | [1 (2 times), 3 (1 time)] |
| 4. | $1 + 4$ | [1 (1 time), 4 (1 time)] |
| 5. | 5 | [5 (1 time)] |
| 6. | $1 + 2 + 2$ | [1 (1 time), 2 (2 times)] |
| 7. | $2 + 3$ | [2 (1 time), 3 (1 time)] |

- ∴ Number of conjugacy classes = 7.
 An element of 1st class = (1)
 An element of 2nd class = (1 2)
 An element of 3rd class = (1 2 3)
 An element of 4th class = (1 2 3 4)
 An element of 5th class = (1 2 3 4 5)
 An element of 6th class = (1 2)(3 4)
 An element of 7th class = (1 2 3)(4 5)

1.13 Give an example of an infinite group in which every element has finite order.

(2013 : 10 Marks)

Solution:

Consider $\langle \frac{Q}{Z}, + \rangle$, i.e., the quotient group of rationals with respect to integers under addition.

Any element of $\frac{Q}{Z}$ is of form $\frac{p}{q} + z$ where $\frac{p}{q} \in Q$ is in lowest form.

$$q\left(\frac{p}{q} + z\right) = \underbrace{\frac{p}{q} + \dots + \frac{p}{q}}_{q \text{ times}} + z = p + z$$

$$= z$$

$$\therefore O\left(\frac{p}{q} + z\right) \leq q$$

So, order of all elements are finite. But there are infinitely elements in $\frac{Q}{Z}$ as there are infinite rationals in $[0, 1)$ and none of them are equal in $\frac{Q}{Z}$.

$$\begin{aligned} \text{And } f[(a+ib)(c+id)] &= f[ac - bd + i(ad+bc)] \\ &= \begin{bmatrix} ac - bd & -(ad+bc) \\ ad+bc & ac - bd \end{bmatrix} = \begin{bmatrix} a & -b \\ b & a \end{bmatrix} \begin{bmatrix} c & -d \\ d & c \end{bmatrix} \\ &= f(a+ib) \cdot f(c+id) \end{aligned}$$

f is one-one

$$a+ib = c+id \Rightarrow a=c \text{ and } b=d \Rightarrow \begin{bmatrix} a & -b \\ b & a \end{bmatrix} = \begin{bmatrix} c & -d \\ d & a \end{bmatrix}$$

$$\Rightarrow f(a+ib) = f(c+id)$$

f is onto

$$\begin{bmatrix} a & -b \\ b & a \end{bmatrix} \in S \begin{bmatrix} a & -b \\ b & a \end{bmatrix} = f(a+ib)$$

∴ f is a one-one onto linear map and so an isomorphism.

1.14 What are the orders of the following permutations in S_{10} .

$$\begin{pmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 & 10 \\ 1 & 8 & 7 & 3 & 10 & 5 & 4 & 2 & 6 & 9 \end{pmatrix} \text{ and } (1\ 2\ 3\ 4\ 5)(6\ 7)$$

(2013 : 10 Marks)

Solution:

Approach : This and the next part uses the fact that order of any permutation written as disjoint cycles in LCM of the order of the cycle. And order of any cycle is its length.

Writing the permutation as product of disjoint cycles.

$$(2\ 8)(3\ 7\ 4)(5\ 10\ 9\ 6)$$

$$\text{Order} = \text{L.C.M.}(2, 3, 4) = 12$$

$$(1\ 2\ 3\ 4\ 5)(6\ 7)$$

$$\text{Order} = \text{L.C.M.}(5, 2) = 10$$

1.15 What is the maximal possible order of an element in S_{10} ? Why? Give an example of that element? How many elements will be there in S_{10} of that order?

(2013 : 13 Marks)

Solution:

Any permutation in S_{10} can be written as product of disjoint cycles. Let n_1, n_2, \dots, n_r be size of these cycles. Then they are also the order of respective cycles.

$$n_1 + n_2 + \dots + n_r = 10$$

and $\text{Order} = \text{L.C.M.}(n_1, n_2, \dots, n_r)$

The order will be maximised if n_i 's are relative primes.

The primes less than 10 are 2, 3, 5, 7.

Taking $n_1 = 3, n_2 = 7$, Order = 21; $n_1 = 2, n_2 = 3, n_3 = 5$, Order = 30.

\therefore The highest order is 30 for cycles of length 2, 3 and 5.

An example : $(1, 2)(3\ 4\ 5)(6\ 7\ 8\ 9\ 10)$

Number of Such Elements :

${}^{10}C_2$ ways of picking 2 elements for 1st cycle and they can be written $(2-1)! = 1$ way.

8C_3 ways of picking 3 elements from rest 8 elements and can be written in $(3-1)! = 2$ ways.

Rest 5 elements can be written in $(5-1)! = 24$ ways.

$$\begin{aligned} \text{Total number of elements} &= {}^{10}C_2 \times 1 + {}^8C_3 \times 2 + 1 \times 24 \\ &= \frac{10 \times 9}{2} + \frac{8 \times 7 \times 6}{3 \times 2} \times 2 + 24 \\ &= 45 + 56 + 24 = 135 \text{ elements.} \end{aligned}$$

1.16 Let G be the set of all real 2×2 matrices $\begin{bmatrix} x & y \\ 0 & z \end{bmatrix}$, where $xz \neq 0$. Show that G group under matrix multiplication. Let N denote the subset $\left\{ \begin{bmatrix} 1 & a \\ 0 & 1 \end{bmatrix} \mid a \in \mathbb{R} \right\}$. Is N a normal subgroup of G ? Justify your answer.

(2014 : 10 Marks)

Solution:

Let $(G, *)$ be an algebraic structure. Where $|*$ implies multiplication between its element in ' G ' is the set of

all 2×2 matrices of type $\begin{bmatrix} x & y \\ 0 & z \end{bmatrix}; xz \neq 0, x, y, z \in R$

$$G = \left\{ \begin{bmatrix} x & y \\ 0 & z \end{bmatrix}; xz \neq 0, x, y, z \in R \right\}$$

(i) Closure Property:

$$\forall \begin{bmatrix} x & y \\ 0 & z \end{bmatrix}, \begin{bmatrix} a & b \\ 0 & c \end{bmatrix} \in G$$

$$x, y, z, a, b \in 1R \tag{... (i)}$$

$$xz \neq a, ac \neq 0 \tag{... (ii)}$$

$$\Rightarrow \begin{bmatrix} x & y \\ 0 & z \end{bmatrix} * \begin{bmatrix} a & b \\ 0 & c \end{bmatrix} = \begin{bmatrix} ax & xb + yc \\ 0 & zc \end{bmatrix} = A$$

$$ax \in 1R$$

$$(ax).(cz) = (ac)(xz) \neq 0 \text{ from (ii)}$$

$$xb + yc \in 1R$$

$$zc \in 1R \text{ from (i)}$$

$$\therefore A \in G \Rightarrow (G, *) \text{ is closed}$$

(ii) Associative Property :

$$\forall \begin{bmatrix} x & y \\ 0 & z \end{bmatrix}, \begin{bmatrix} a & b \\ 0 & c \end{bmatrix}, \begin{bmatrix} p & a \\ 0 & r \end{bmatrix} \in G$$

$$x, y, z, a, b, c, p, r \in 1R$$

$$xz \neq 0, ac \neq 0, Pr \neq 0$$

$$\Rightarrow \left(\begin{bmatrix} x & y \\ 0 & z \end{bmatrix} * \begin{bmatrix} a & b \\ 0 & c \end{bmatrix} \right) * \begin{bmatrix} p & a \\ 0 & r \end{bmatrix} = \begin{bmatrix} ax & bx + yc \\ 0 & zc \end{bmatrix} * \begin{bmatrix} p & a \\ 0 & r \end{bmatrix}$$

$$= \begin{bmatrix} apx & aqx + bxr + ycr \\ 0 & zcr \end{bmatrix} \tag{... (iii)}$$

$$\begin{bmatrix} x & y \\ 0 & z \end{bmatrix} * \left(\begin{bmatrix} a & b \\ 0 & c \end{bmatrix} * \begin{bmatrix} p & a \\ 0 & r \end{bmatrix} \right) = \begin{bmatrix} apx & aqx + bxr + ycr \\ 0 & zcr \end{bmatrix} \tag{... (iv)}$$

Here, (iii) = (iv)

$$\text{i.e. } \left(\begin{bmatrix} x & y \\ 0 & z \end{bmatrix} * \begin{bmatrix} a & b \\ 0 & c \end{bmatrix} \right) * \begin{bmatrix} p & a \\ 0 & r \end{bmatrix} = \begin{bmatrix} x & y \\ 0 & z \end{bmatrix} * \left(\begin{bmatrix} a & b \\ 0 & c \end{bmatrix} * \begin{bmatrix} p & a \\ 0 & r \end{bmatrix} \right)$$

∴ (G, x) satisfy associative property.

(iii) Existence of Left Identity :

$$\text{Let } \begin{bmatrix} x & y \\ 0 & z \end{bmatrix} \in G, \text{ where } xz \neq 0, x, y, z \in 1R \tag{... (v)}$$

$$\text{Let } \begin{bmatrix} a & b \\ 0 & c \end{bmatrix} \in G, \text{ where } ac \neq 0, a, b, c \in 1R \tag{... (vi)}$$

$$\begin{bmatrix} a & b \\ 0 & c \end{bmatrix} * \begin{bmatrix} x & y \\ 0 & z \end{bmatrix} = \begin{bmatrix} x & y \\ 0 & z \end{bmatrix}$$

$$\begin{bmatrix} ax & ay + bz \\ 0 & cz \end{bmatrix} = \begin{bmatrix} x & y \\ 0 & z \end{bmatrix}$$

equating its element,

$$cz = z \Rightarrow z(c - 1) = 0$$

$$\text{From (vi), } ac \neq 0 \Rightarrow a \neq 0, c \neq 0$$

$$\text{From (v) } xz \neq 0 \Rightarrow x \neq 0, z \neq 0$$

$$\therefore c - 1 = 0 \Rightarrow c = 1$$

$$ax = x \Rightarrow x(1 - a) = 0$$