



**POSTAL
BOOK PACKAGE**

2025

CONTENTS

**MECHANICAL
ENGINEERING**

Objective Practice Sets

Renewable Sources of Energy

1. Solar Radiation 2 - 8
2. Solar Thermal Energy Collection 9 - 12
3. Solar Thermal Energy Storage and Applications 13 - 17
4. Indirect Sources of Solar Energy 18 - 20
5. Renewable Energy Sources Other than Solar Energy .. 21 - 23

Solar Radiation

- Q.1** Estimate the surface temperature of the sun if extraterrestrial radiation flux received are 1367 W/m^2 . The mean distance between the Sun and Earth is $1.496 \times 10^8 \text{ km}$ and radius of sun is $6.95 \times 10^5 \text{ km}$. You can assume Sun as blackbody.
- (a) 5780 K (b) 5880 K
(c) 4088 K (d) 6180 K
- Q.2** A scientist is carrying out experiments on solar plate collectors in an observatory located at 78°N , 38°E . Due to malfunction of instruments, experimental results are not available. Find out theoretical value of extraterrestrial radiation on 21st June for the given location.
- (a) 1367 W/m^2 (b) 1441 W/m^2
(c) 1353 W/m^2 (d) 1323 W/m^2
- Q.3** What is the orientation angle of Earth and solar flux received outside the atmosphere of Earth on 14th November, 2016?
- (a) 16.47° , 1398 W/m^2
(b) -19.15° , 1116 W/m^2
(c) 16.47° , 1116 W/m^2
(d) -19.15° , 1398 W/m^2
- Q.4** The hour angle made by Earth at a location in New Delhi ($25^\circ35'\text{N}$, $77^\circ12'\text{E}$) at 1200 h (LAT) will be _____.
- (a) -7.5° (b) 7.5°
(c) 15° (d) zero
- Q.5** The atmosphere of earth is
- (a) opaque for short-wavelength but transparent for long-wavelength radiation.
(b) transparent for short-wavelength radiation but opaque for long-wavelength radiation.
(c) transparent for both short and long-wavelength radiation.
(d) opaque for both short and long-wavelength radiation.
- Q.6** In solar radiation, the term 'albedo' is used to quantify:
- (a) amount of radiation reflected back to the space
(b) amount of diffuse radiation received on the surface of earth
(c) amount of air-mass ratio for maximum radiation
(d) amount of global radiation received on earth
- Q.7** Find out hour angle at a location in Shillong ($25^\circ34'\text{N}$, $91^\circ56'\text{E}$) on 23rd May 2006 at 1000 h (IST). Standard time in India is based on 82.5°E .
- (a) -30° (b) -22.5°
(c) -20.5° (d) -15°
- Q.8** Find out local apparent time in Jodhpur ($26^\circ18'\text{N}$, $73^\circ01'\text{E}$)
- (a) 1008 h (b) 0.852 h
(c) 0930 h (d) 0922 h
- Q.9** A flat plate collector is located in New Delhi ($28^\circ35'\text{N}$, $77^\circ12'\text{E}$) and tilted at an angle 36° with horizontal. The collector is facing due south. Find out air-mass ratio on 30th June at 0945 (LAT).
- (a) 1.24 (b) 1.70
(c) 1.16 (d) 1.96
- Q.10** The term 'turbidity' indicates:
- (a) Turbulence intensity in the atmosphere
(b) Haziness caused by invisible particles
(c) Turbulence in the core region of sun
(d) Air velocity in the upper atmosphere
- Q.11** Solar beam radiation falling on a particular in Bulgaria are $800 \text{ W/m}^2\text{-hr}$. If diffused radiation are 10% of the beam radiation, what would be global radiation?
- (a) $720 \text{ W/m}^2\text{-hr}$ (b) $65 \text{ W/m}^2\text{-hr}$
(c) $880 \text{ W/m}^2\text{-hr}$ (d) $100 \text{ W/m}^2\text{-hr}$
- Q.12** Solar constant is obtained from spectral distribution of extraterrestrial radiation. It is the value corresponding to:
- (a) maximum wavelength in the spectra
(b) minimum wavelength in the spectra

- (c) mean wavelength in the spectra
- (d) area weighted average over the range of wavelengths in spectra

Q.13 A flat plate collector is used for water heating in Jammu (32.72°N, 74.86°E). The collector is inclined at an angle 20° and facing due South. Find out solar beam incident angle (θ) and solar altitude at 1100 h (local apparent time) on 8th August.

- (a) 15.5°, 21.57° (b) 15.5°, 68.43°
- (c) 21.57°, 74.5° (d) 15.5°, 74.5°

Q.14 On a hazy day in Bhopal (23.26°N, 77.41°E) at 1030 h (LAT) air-mass ratio is 1.84. Find out solar azimuth angle on 20th Jan 2008 for the given data.

- (a) 57° (b) 43°
- (c) 77° (d) 23°

Q.15 A flat plate collector located in Hyderabad (17.39° N, 78.49° E), is used to supply dry air. Find out sunrise hour angle on 23rd March at 730 h (local apparent time). Also find out maximum sunshine hours for the given details.

- (a) 90°, 12 h (b) 450°, 6h
- (c) 45°, 12 h (d) 90°, 6 h

Q.16 A concentrated collector is used to heat up brine. The concentrator is located in Lucknow (26.85°N, 80.95°E) and rotated about horizontal east-west axis. The collector is adjusted once a day to track sun. Find out incidence angle on 2nd Feb at 1030 h (LAT) and slope of the surface for the given day respectively.

- (a) 21.5°, 44° (b) 44°, 21.5°
- (c) 27.5°, 36° (d) 36°, 27.5°

Q.17 A solar concentrated collector unit is installed in Bhubaneswar (20.30°N, 85.82°E). Find out angle of incidence and solar azimuth angle on 31st Jan, 1400 h, if it is rotated about east-west for continuous adjustment.

- (a) 28°, 39° (b) 28°, 48°
- (c) 39°, 28° (d) 39°, 48°

Q.18 A space rover receives power from solar radiation. Horizontal Flat plate collectors of 8 m² area are used for this purpose. The flaps of collector will be opened after crossing atmospheric boundary

layer. Space rover will left Earth's atmosphere on 1st May. Find out extraterrestrial radiation falling on rover on a particular time when solar azimuth angle is 60° for the data given

- (a) 1367 W (b) 1036 W
- (c) 5468 W (d) 5380 W

Q.19 Find out the solar radiation on horizontal surface in the absence of the atmosphere at Gurugram (28.46°N, 77.02°E) on 15th April between 0900 to 1000 h.

- (a) 20.85 MJ/m² (b) 3.73 MJ/m²
- (c) 1036 KJ/m² (d) 1202 KJ/m²

Q.20 The monthly average daily extraterrestrial radiation at a location (43°5, 80° ω) is 3.5 MJ/m² on 23rd September. Due to same haziness, the clarity index is 0.74. Find out the monthly average daily global radiation if prediction model of page i.e.

$$\frac{\bar{H}_g}{H_0} = a + b \left(\frac{\bar{S}}{\bar{S}_{\max}} \right)$$

is applicable for estimating solar radiation. The value of regression constants are 0.31 and 0.58 respectively. Also find out declination angle of the Earth if observed sunshine hours for the given location are 9 h.

- (a) 2.59 MJ/m², 11.26°
- (b) 4.73 MJ/m², 11.26°
- (c) 2.59 MJ/m², 14.26°
- (d) 2.59 MJ/m², 1.126°

Q.21 The latitude angle is zero at

- (a) Southern Hemisphere
- (b) Northern Hemisphere
- (c) Equator
- (d) All of the above

Q.22 The extraterrestrial solar intensity flux is dependent on:

- (a) Height from the surface of earth
- (b) Atmospheric conditions
- (c) Locations of the observer
- (d) Day of the year

Q.23 Which of the following angle varies seasonally due to the tilt of the earth on its axis and rotations of the earth around the sun?

- (a) Tilt angle (b) Altitude angle
- (c) Hour angle (d) Declination angle

- Q.24** Which of the following statements is correct in regard of solar Zenith angle?
 (a) It is zero at the time of sunrise
 (b) It is zero at the solar noon
 (c) It is zero at the time of sun fall
 (d) It is maximum at solar noon
- Q.25** Which of the following instruments is used to measure direct/beam radiation?
 (a) Pyrheliometer
 (b) Pyranometer
 (c) Anemometer
 (d) Albedometer
- Q.26** The purpose of shading ring which is being used in pyranometer is/are
 (a) to avoid overheating of the sensor.
 (b) to measure diffuse radiation.
 (c) to measure day length i.e. sunfall-sunrise hours.
 (d) to measure direct radiation.
- Q.27** Solar flux are reported in 'Langley' sometimes, which is the unit of radiation adopted after the name of Samuel Langley. 1 Langley is equal to _____.
 (a) 1 cal (b) 1 kcal
 (c) 1 cal/cm² (d) 1kcal/cm²



Answers Solar Radiation

1. (a) 2. (d) 3. (d) 4. (d) 5. (b) 6. (a) 7. (c) 8. (d) 9. (c)
 10. (b) 11. (c) 12. (d) 13. (b) 14. (b) 15. (a) 16. (b) 17. (b) 18. (d)
 19. (b) 20. (d) 21. (c) 22. (d) 23. (d) 24. (b) 25. (a) 26. (b) 27. (c)

Explanations Solar Radiation

1. (a)

The total emitted radiation by the surface of the Sun will be given by:

$$\sigma T_s^4 (4\pi R_s^2)$$

These radiation are received by a sphere having radius equal to mean distance between the Sun and the Earth.

i.e. $I_{sc} \times (4\pi R_m^2)$

Hence $\sigma T_s^4 (4\pi R_s^2) = I_{sc} \times (4\pi R_m^2)$

$$\Rightarrow \sigma T_s^4 = 1367 \times \left(\frac{1.496 \times 10^{11}}{6.95 \times 10^8} \right)^2$$

$$\Rightarrow \begin{aligned} T_s &= (11.17 \times 10^{14})^{1/4} \\ T_s &= 5781.22 \text{ K} \end{aligned}$$

2. (d)

Extraterrestrial solar flux (I'_{sc}) is given by:

$$I'_{sc} = I_{sc} \left[1 + 0.033 \cos \frac{360n}{365} \right]$$

Solar constant,

$$I_{sc} = 1367 \text{ W/m}^2$$

For 21st June, $n = 31 + 28 + 31 + 30 + 31 + 21$
 $= 172$

$$\begin{aligned} \therefore I'_{sc} &= 1367 \left[1 + 0.33 \cos \frac{360 \times 172}{365} \right] \\ &= 1322.62 \text{ W/m}^2 \end{aligned}$$

Note: Location data and 21st June is given to create confusion. On 21st Jun, Earth's declination angle will be 23.45° i.e. default value. One should not select default value of I_{sc} i.e. 1367 W/m² for the given data.

3. (d)

The variation of Earth's orientation angle is given by cooper's relation:

$$\delta = 23.45 \sin \left[\frac{360}{365} (284 + n) \right]$$

On 14th November

$$\begin{aligned} n &= 31 + 29 + 31 + 30 + 31 + 30 \\ &\quad + 31 + 31 + 30 + 31 + 14 = 319 \end{aligned}$$

$$\delta = 23.45 \sin \left[\frac{360}{365} (284 + 319) \right]$$

$$= -19.15^\circ$$

Flux received outside of the atmosphere of Earth i.e. extraterrestrial flux (I'_{sc}) is given by: